

Dose-Response: An International Journal

Volume 8 | Issue 1

Article 15

3-2010

HORMETIC EFFECTS OF REACTIVE OXYGEN SPECIES BY EXERCISE: A VIEW FROM ANIMAL STUDIES FOR SUCCESSFUL AGING IN HUMAN

Sataro Goto

Juntendo University Graduate School, Inbagun, Japan, and Semmelweis University, Budapest, Hungary

Zsolt Radák

Semmelweis University, Budapest, Hungary

Follow this and additional works at: https://scholarworks.umass.edu/dose_response

Recommended Citation

Goto, Sataro and Radák, Zsolt (2010) "HORMETIC EFFECTS OF REACTIVE OXYGEN SPECIES BY EXERCISE: A VIEW FROM ANIMAL STUDIES FOR SUCCESSFUL AGING IN HUMAN," *Dose-Response: An International Journal*: Vol. 8 : Iss. 1 , Article 15.

Available at: https://scholarworks.umass.edu/dose_response/vol8/iss1/15

This Article is brought to you for free and open access by ScholarWorks@UMass Amherst. It has been accepted for inclusion in Dose-Response: An International Journal by an authorized editor of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

HORMETIC EFFECTS OF REACTIVE OXYGEN SPECIES BY EXERCISE: A VIEW FROM ANIMAL STUDIES FOR SUCCESSFUL AGING IN HUMAN

Sataro Goto □ Juntendo University Graduate School, Institute of Health and Sports Science & Medicine, Inbagan, Japan, and Faculty of Physical Education and Sport Science, Semmelweis University, Budapest, Hungary

Zsolt Radák □ Faculty of Physical Education and Sport Science, Semmelweis University, Budapest, Hungary

□ Numerous anti-aging measures have been proposed to cope with age-associated decline of physiological functions and/or onset of diseases, mostly based on free radical (or oxidative stress) theory of aging, though no robust scientific data have been reported to extend human healthspan. This is due to dual (harmful as well as essential) roles of reactive oxygen species (ROS) to a body. Regular moderate exercise provides benefits upregulating defense against oxidative stress in good balance between the opposing dual roles. Sources of ROS in exercise appear to be not only mitochondria as often claimed but also enzymatic reactions catalyzed by NADPH oxidase and other oxidases. It may, therefore, be possible to mimic this aspect of exercise to promote the defense for healthspan extension by other means such as modest alcohol consumption that could upregulate activity of enzymes against oxidative stress.

Keywords: exercise, hormesis, reactive oxygen species, anti-oxidant, healthspan

INTRODUCTION

Despite remarkable prolongation of lifespan in developed countries in the last several decades, a considerable proportion of the elderly population is frail and vulnerable to physical and mental disorders that impairs quality of their later lives. Numerous anti-aging measures have been proposed to ameliorate and/or retard age-associated decline of physiological functions and/or onset of diseases. A most popular working hypothesis on mechanisms of aging, free-radical theory of aging/oxidative stress theory of aging (Harman 2009), has been the basis of major anti-aging strategies, although the theory has been criticized as being not definitely proven (Muller et al. 2007). In accordance with such claims, anti-oxidants including vitamin C and E have been used by people with little or no health promoting effects except in obvious disease conditions in an attempt to reduce damage due to reactive oxygen species (ROS) that may drive aging and cause diseases. These vitamins and anti-oxidants such as catechins and other polyphenols can even be detrimental to anti-

Address correspondence to Sataro Goto, Juntendo University Graduate School, Institute of Health and Sports Science & Medicine, Hiragagakuendai 1-1, Inbamura, Inbagan, Chiba, 270-1695 Japan; Phone: +81-476-98-1001, Fax: +81-476-98-1030, Email: gotosataro@sakura.juntendo.ac.jp

Exercise hormesis for healthspan extension

mals and human (Bjelakovic *et al.* 2008). These facts, however, are not necessarily contradictory to the oxidative theory of aging in view of possible hormetic effects of ROS.

OPPOSING ROLES OF ROS AND CRITIC OF USING ANTI-OXIDANTS TO REDUCE OXIDATIVE STRESS

ROS are generated in a variety of cellular processes. Superoxide anion radicals are byproducts of electron transport system in mitochondria and endoplasmic reticulum. Chemical reactions catalyzed by oxidases such as NADPH oxidase, xanthine oxidase and monoamine oxidase produce superoxide and/or hydrogen peroxide as normal and inevitable products. Highly reactive hydroxyl radicals generated from hydrogen peroxide by transition metal catalyzed Fenton reaction can damage proteins, nucleic acids and membrane phospholipids. On the other hand, it has been clearly demonstrated that superoxide and/or hydrogen peroxide have essential physiological roles in cellular functions such as signal transduction, cell proliferation and transcription regulation (Droge 2002). Because of such opposing roles of ROS, anti-oxidants often claimed to reduce oxidative stress have been criticized as having no significant effects or potentially detrimental consequences.

EXERCISE HORMESIS AND POSSIBLE HUMAN HEALTHSPAN EXTENSION BY MODERATE OXIDATIVE STRESS

Extensive physical activities in unprepared body can cause oxidative damage of nucleic acids, proteins and membrane lipids in various tissues (Davies *et al.* 1982) due to massive generation of ROS. ROS are generated during and/or after exercise in mitochondrial respiratory chain, and reactions catalyzed by monoamine oxidase (Cadenas and Davies 2000), NADPH oxidase or myeloperoxidase in neutrophils mobilized or infiltrated in the skeletal muscle (Peake and Suzuki, 2004), xanthine oxidase (Viña *et al.* 2000) as well as from endothelial cells in vascular systems (Rush and Aultman 2008). It should be noted that ROS generation from respiring mitochondria is claimed to be one or two orders of magnitude lower than reported (*i.e.*, 2 to 3 % of oxygen consumed) in the skeletal muscle and liver under conditions that mimic physiological situations (St-Pierre *et al.* 2002). Major sites of ROS generation are not only mitochondria but also other sources such as xanthine oxidase and NADPH oxidase catalyzed reactions (Bonetto *et al.* 2009). Hypoxia-induced generation of ROS can also be an important source (Clanton 2007). Due to massive consumption of oxygen in the skeletal muscle during exercise, transient lowering of oxygen concentration may result in other tissues, thus causing temporary local hypoxia. It is, therefore, likely that modest increase of ROS by exercise can induce defense enzymes and other proteins

against oxidative stress systemically in cells and tissues unprepared for excessive generation of ROS. Such situations would lead to adaptation to stronger stress that may be encountered in future.

Indeed, we have shown that regular swimming exercise in young and middle-aged rats can reduce protein carbonyls in the brain that are accompanied with upregulation of the activity of proteasome for degradation of the oxidatively modified proteins (Radák et al. 2001). Similarly, regular treadmill running reduced oxidative damage to DNA in the skeletal muscle (Radák et al. 1999) and liver (Nakamoto et al. 2007) of aged rats, with the increase in the repair enzyme activity (OGG1). Higher protection against oxidative stress was also observed in the myocardium after regular swimming exercise (Radák et al. 2000). It was also demonstrated that regular treadmill running apparently attenuated age-associated increase in oxidative stress by reducing NF- κ B activation with increased level of reduced form of glutathione in the liver of aged rats (Radák et al. 2004). Others have reported upregulation of antioxidant enzymes by exercise (e.g., Ji et al. 2006). Thus, mild oxidative stress induced by exercise appears to be able to reduce oxidative damage by upregulating antioxidant systems, collectively termed as primary (e.g. glutathione), secondary (e.g. glutathione peroxidase) and tertiary (e.g. OGG1 and proteasome) defense system. As such, we have argued that dual (harmful and beneficial) effects of ROS generated by exercise can be a form of hormesis (Radák et al. 2005, 2008; Ji et al. 2008). Supporting this argument, we have shown that hydrogen peroxide (oxidant) or N-tert butyl-alpha-phenyl nitron (anti-oxidant) injection modulates neurotrophin levels in the spinal cord in rats, apparently via alteration of redox state (Siamilis et al. 2009).

In agreement with the idea of exercise hormesis caused by ROS in animals, human studies of antioxidant supplementation have been reported to attenuate beneficial effects of exercise. Khassaf et al. (2003) found that long term vitamin C supplementation attenuated adaptive response of exercise to oxidants in human lymphocytes. Fischer et al. (2004) reported that supplementation of vitamin C and E blunted exercise induced IL-6 release from contracting human skeletal muscle. More recently, Ristow et al. (2009) demonstrated that health promoting effects of physical exercise such as upregulation of glucose infusion rate, plasma adiponectin level and mRNA of anti-oxidant enzymes in human skeletal muscle are abolished by administration of vitamin C and E.

Thus, ROS generated by physical activity appears to upregulate defense against oxidative stress and other beneficial effects in both rodents and humans. It is, therefore, likely that mild oxidative stress is useful in retarding physiological decline and reducing risk of diseases associated with aging whether it is caused by exercise or perhaps by other means (see below).

CONCLUSION AND PERSPECTIVES

Obviously, too much physical activity is harmful due to massive generation of ROS, if a body is not conditioned to minimize damage. Modest and regular exercise, however, can induce cellular defense against stronger oxidative stress, promoting health and reducing risk of diseases. Importantly, such hormetic adaptation can occur even at old ages at least in model animals (Goto et al.2004; Cui et al.2009). It has not been demonstrated whether it is true in older humans as has been shown in young individuals as cited above. It is interesting to point out that alcohol consumption, if modest, promotes health despite it causes oxidative stress. Notably, prior intake of alcohol can attenuate ischemic/reperfusion damage in the brain in model animals due to ROS generation in NADPH oxidase catalyzed reaction (Wang et al. 2007). Thus, moderate alcohol intake can have hormetic effects similar to exercise via ROS generation. Lipopolysaccharide (LPS), a bacterial endotoxin that induces inflammation, activates NADPH oxidase dependent ROS production from neutrophils (Brandes et al.1999). LPS might also be viewed as an agent that could cause beneficial outcomes if doses would not be high, as in the case of non-severe bacterial infection, thereby possibly promoting anti-oxidant defense. It would be interesting to explore means for mild oxidative stress such as those suggested in the effect of polyphenols as prooxidants that can induce defense against stronger harmful stress, rather than reducing oxidative stress by their antioxidant activity *per se* (Stevenson and Hurst 2007).

REFERENCES

- Bonetto A, Penna F, Muscaritoli M, Minero VG, Fanelli FR, Baccino FM and Costelli P. 2009. Are antioxidants useful for treating skeletal muscle atrophy? *Free Radic Biol Med.* 47: 906-916
- Bjelakovic G, Nikolova D, Simonetti RG and Gluud C. 2008. Systematic review: primary and secondary prevention of gastrointestinal cancers with antioxidant supplements. *Aliment Pharmacol Ther.* 28: 689-703
- Brandes RP, Koddenberg G, Gwinner W, Kim D, Kruse HJ, Busse R and Mügge A. 1999. Role of increased production of superoxide anions by NAD(P)H oxidase and xanthine oxidase in prolonged endotoxemia. *Hypertension.* 33:1243-1249
- Cadenas E and Davies KJ. 2000. Mitochondrial free radical generation, oxidative stress, and aging. *Free Radic Biol Med.* 29: 222-230
- Clanton TL. 2007. Hypoxia-induced reactive oxygen species formation in skeletal muscle. *J Appl Physiol.* 102: 2379-2388
- Cui L, Hofer T, Rani A, Leeuwenburgh C and Foster TC. 2009. Comparison of lifelong and late life exercise on oxidative stress in the cerebellum. *Neurobiol Aging.* 30: 903-909
- Davies KJ, Quintanilha AT, Brooks GA and Packer L. 1982. Free radicals and tissue damage produced by exercise. *Biochem Biophys Res Commun.* 107: 1198-205
- Droge W. 2002. Free radicals in the physiological control of cell function. *Physiol Rev.* 82: 47-95
- Fischer CP, Hiscock NJ, Penkowa M, Basu S, Vessby B, Kallner A, Sjöberg LB and Pedersen BK. 2004. Supplementation with vitamins C and E inhibits the release of interleukin-6 from contracting human skeletal muscle. *J Physiol.* 558: 633-645
- Goto S, Radák Z, Nyakas C, Chung HY, Naito H, Takahashi R, Nakamoto H and Abe R. 2004. Regular exercise: An effective means to reduce oxidative stress in old rats. *Ann NY Acad Sci* 1019: 471-474

- Harman D. 2009. Origin and evolution of the free radical theory of aging: a brief personal history, 1954-2009. *Biogerontology*. DOI 10.1007/s10522-009-9234-2 (moved and DOI No added)
- Ji LL, Gomez-Cabrera MC and Vina J. 2006. Exercise and hormesis: activation of cellular antioxidant signaling pathway. *Ann NY Acad Sci*. 1067: 425-435
- Ji LL, Radák Z and Goto S. 2008. Hormesis and Exercise: How the cell copes with oxidative stress. *Am J Pharmacol Toxicol* 3: 41-55
- Khassaf M, McArdle A, Esanu C, Vasilaki A, McArdle F, Griffiths RD, Brodie DA and Jackson MJ. 2003. Effect of vitamin C supplements on antioxidant defence and stress proteins in human lymphocytes and skeletal muscle. *J Physiol*. 549: 645-652
- Muller FL, Lustgarten MS, Jang Y, Richardson A and Van Remmen H. 2007. Trends in oxidative aging theories. *Free Radic Biol Med*. 43:477-503
- Nakamoto H, Kaneko T, Tahara S, Hayashi E, Naito H, Radák Z and Goto S. 2007. Regular exercise reduces 8-oxodG in the nuclear and mitochondrial DNA and modulates the DNA repair activity in the liver of old rats. *Exp Gerontol* 42: 287-295
- Peake J and Suzuki K. 2004. Neutrophil activation, antioxidant supplements and exercise-induced oxidative stress. *Exerc Immunol Rev*. 10: 129-141.
- Radák Z, Kaneko T, Tahara S, Nakamoto H, Ohono H, Sasvari M, Nyakas C and Goto S. 1999. The effect of exercise training on oxidative damage of lipids, proteins and DNA in rat skeletal muscle: Evidence for beneficial outcome. *Free Radic Biol Med* 27: 69-74
- Radák Z, Sasvari M, Nyakas C, Pucso J, Nakamoto H and Goto S. 2000. Exercise preconditioning against hydrogen peroxide-induced oxidative damage in proteins of rat myocardium. *Arch Biochem Biophys* 376: 248-251
- Radák Z, Kaneko T, Tahara S, Nakamoto H, Msasvai M, Nyakas C and Goto S. 2001. Regular exercise improves cognitive function and decreases oxidative damage to proteins in rat brain. *Neurochem Int* 38: 17-23
- Radák Z, Chung HY, Naito H, Takahashi R, Jung KJ, Kim HJ and Goto S. 2004. Age-associated increase in oxidative stress and nuclear transcription factor NF- κ B activation are attenuated in rat liver by regular exercise. *FASEB J* 18: 749-750
- Radák Z, Chung HY and Goto S. 2005. Opinion: Exercise and hormesis: Oxidative stress-related adaptation for successful aging. *Biogerontology* 6: 71-75
- Radák Z, Chung HY, Koltai E, Taylor AW and Goto S. 2008. Exercise, oxidative stress and hormesis. *Ageing Res Rev* 7: 34-42
- Ristow M, Zarse K, Oberbach A, Klötting N, Birringer M, Kiehntopf M, Stumvoll M, Kahn CRF and Blüher M. 2009. Antioxidants prevent health-promoting effects of physical exercise in humans. *Proc Natl Acad Sci U S A*. 106: 8665-8670
- Rush JW and Aultman CD. 2008. Vascular biology of angiotensin and the impact of physical activity. *Appl Physiol Nutr Metab*. 33: 162-172 (moved)
- Siamilis S, Jakus J, Nyakas C, Costa A, Mihalik B, Falus A and Radák Z. 2009. The effect of exercise and oxidant-antioxidant intervention on the levels of neurotrophins and free radicals in spinal cord of rats. *Spinal Cord*. 47: 453-457
- Stevenson DE and Hurst RD. 2007. Polyphenolic phytochemicals—just antioxidants or much more? *Cell Mol Life Sci*. 64: 2900-2916
- St-Pierre J, Buckingham JA, Roebuck SJ and Brand MD. 2002. Topology of superoxide production from different sites in the mitochondrial electron transport chain. *J Biol Chem*. 277: 44784-44790
- Viña J, Gimeno A, Sastre J, Desco C, Asensi M, Pallardó FV, Cuesta A, Ferrero JA, Terada LS and Repine JE. 2000. Mechanism of free radical production in exhaustive exercise in humans and rats; role of xanthine oxidase and protection by allopurinol. *IUBMB Life*. 49: 539-544
- Wang Q, Sun AY, Simonyi A, Kalogeris TJ, Miller DK, Sun GY and Korthuis RJ. 2007. Ethanol preconditioning protects against ischemia/reperfusion-induced brain damage: role of NADPH oxidase-derived ROS. *Free Radic Biol Med*. 43: 1048-1060